Rita, Rice and Salt Water: An Overview of the Aftermath of Hurricane Rita in the Southwest Louisiana Rice and Sugarcane Growing Area

Johnny Saichuk, Rice Specialist LSU AgCenter



LSU AgCenter Scientists Involved in Salt Studies

Dr. Brooks Blanche

Dr. Jason Bond

Dr. Gary Breitenbeck

Mr. Tom Hymel

Dr. Ben Legendre

Dr. Steve Linscombe

Mr. J Stevens

Dr. Ed Twidwell

Dr. Howard Viator


























Proposed Ranking Rules of Salt Effects on Rice Production

Rank	Description	Salts (ppm)	SAR	Effects
1	None	<500	<4	No effect on yield
2	Mild	500- 1000	<5	Little to no effect on yields
3	Moderate	1000- 2000	<6	Some yield reduction possible
4	Severe	2000- 6000	6-15	Substantial yield reduction
5	V. Severe	>6000	>15	Catastrophic failure


Class	Salts ppm		SAR	
None	< 500	and	<4	
Mild	500-1000	or	<4	
Moderate	1000-2000	or	<6	
Severe	2000-6000	or	<15	
Very Severe	>6000	Of	>15	




Class	Salts ppm	- 10	SAR	
None	< 500	and	<4	
Mild	500-1000	01	<4	
Moderate	1000-2000	01	<6	
Severe	2000-6000	01	<15	
Very Severe	>6000	or	>15	



Class	Salts ppm		SAR	
None	< 500	and	<4	
Mild	500-1000	or	<4	
Moderate	1000-2000	O1'	<6	
Severe	2000-6000	01	<15	
Very Severe	>6000	Of	>15	



Class	Salts ppm	W 10 10 10	SAR	
None	< 500	and	<4	
Mild	500-1000	Of	<4	
Moderate	1000-2000	or	<6	
Severe	2000-6000	Of	<15	
Very Severe	>6000	or	>15	



Class	Salts ppm		SAR	
None	< 500	and	<4	
Mild	500-1000	or	<4	
Moderate	1000-2000	Of	<6	
Severe	2000-6000	01	<15	
Very Severe	>6000	or	>15	



Class	Salts ppm	+50	SAR	
None	< 500	and	<4	
Mild	500-1000	Of	<4	
Moderate	1000-2000	Of	<6	
Severe	2000-6000	Of	<15	
Very Severe	>6000	01	>15	




Distribution of Sampled Sites Based on Sampling Depth and Anticipated Effects of Salt

Rank	Description	0-3"	3-6"	6-12"	0-6"	0-12"
1	None	31%	41%	47%	35%	41%
2	Mild	11%	20%	19%	19%	15%
3	Moderate	7%	13%	16%	10%	16%
4	Severe	32%	23%	15%	30%	25%
5	V. Severe	17%	3%	3%	6%	3%

Potential Salinity Problems for Sugarcane


- Soil sampling found from 268 to >6,000 ppm of salt in the surface 3 inches where sugarcane fields were exposed to flood waters of Hurricanes Katrina and Rita
- Approximately 40,000 acres of sugarcane area flooded with water containing salt
- Sugarcane is moderately sensitive to salt with a threshold for yield reduction of approximately 1,100 ppm
- Saltiest areas were just north of Hwy 14 and at Burns Point
- Salt content higher in Vermilion Parish and became brackish more eastward


(Viator et. al. 2006)



























Water Vapor

Zone of Salt accumulation

Water movement by capillarity

Current Recommendations

- Do nothing
- Pray for rain
- Sample same areas to detect any changes

Johnny Saichuk 1373 Caffey Rd. Rayne, LA 70578

Office: 337-788-7547

Cell: 337-849-6253

e-mail:

jsaichuk@agcenter.lsu.edu web:lsuagcenter.com/en/crops_ livestock/crops/rice